

**For Reference**

---

**NOT TO BE TAKEN FROM THIS ROOM**



# For Reference

---

NOT TO BE TAKEN FROM THIS ROOM

Ex LIBRIS  
UNIVERSITATIS  
ALBERTAENSIS







Digitized by the Internet Archive  
in 2019 with funding from  
University of Alberta Libraries

<https://archive.org/details/Jorstad1967>





ON A PARTICULAR CLASS OF SIGMA-MONOGENIC FUNCTIONS

by



RALPH JORSTAD

A THESIS

SUBMITTED TO THE FACULTY OF GRADUATE STUDIES

IN PARTIAL FULFILLMENT OF THE REQUIREMENTS FOR THE DEGREE

OF MASTER OF SCIENCE

DEPARTMENT OF MATHEMATICS

UNIVERSITY OF ALBERTA

EDMONTON, ALBERTA

JUNE, 1967





UNIVERSITY OF ALBERTA

FACULTY OF GRADUATE STUDIES

The undersigned certify that they have read and recommend to the Faculty of Graduate Studies for acceptance, a thesis entitled "On a Particular Class of Sigma-monogenic Functions", submitted by RALPH JORSTAD in partial fulfillment of the requirements for the degree of Master of Science.



## ABSTRACT

This thesis is devoted to a study of a particular class of sigma-monogenic functions, namely those whose real and imaginary parts,  $u$  and  $v$ , satisfy

$$\begin{aligned} e^{-cx} u_x &= v_y \\ e^{-cx} u_y &= -v_x \end{aligned} .$$

The idea of a sigma-monogenic function is explained in Chapter I, and by solving Cauchy problems we show how to express our  $\Sigma$ -monogenic functions in terms of analytic functions. The concept of correspondence between  $\Sigma$ -monogenic and analytic functions is also defined here. In Chapter II, the formal powers  $Z^{(n)}(z_0; z)$  are given and bounds on  $|a \cdot Z^{(n)}(z_0; z)|$  are determined. Then in Chapter III, bounds on  $f^{[n]}(z)$ , the derivatives of a  $\Sigma$ -monogenic function, analogous to Cauchy's inequality, are derived by solving a Dirichlet problem in a circle. We are thus able to show that

$$f(z) = \sum_{n=0}^{\infty} \frac{f^{[n]}(z_0)}{n!} Z^{(n)}(z_0; z) \quad \text{for all } |z - z_0| < a \quad \text{provided } f(z) \text{ is}$$

$\Sigma$ -monogenic in that region.

In the Appendix, an alternate proof is given for our formulae expressing a  $\Sigma$ -monogenic function in terms of an analytic function.





#### ACKNOWLEDGEMENTS

I would like to express my sincere gratitude to Dr. J. Sanders for his careful and considerate guidance throughout the preparation of this thesis.



# TABLE OF CONTENTS

	<u>Page</u>
ABSTRACT . . . . .	(i)
ACKNOWLEDGEMENTS . . . . .	(ii)
CHAPTER I: INTRODUCTION . . . . .	1
CHAPTER II: THE POWERS $a \cdot z^{(n)}(z_0; z)$ AND THEIR BOUNDS . . .	13
CHAPTER III: ESTIMATES ON THE DERIVATIVES OF $f(z)$ AND THE GENERALIZED TAYLOR'S THEOREM . . . . .	22
BIBLIOGRAPHY . . . . .	38
APPENDIX . . . . .	39





# CHAPTER I

## INTRODUCTION

In 1944, Bers and Gelbart [1] considered the system of equations

$$\begin{aligned}\sigma_1(x) u_x &= \tau_1(y) v_y \\ \sigma_2(x) u_y &= -\tau_2(y) v_x\end{aligned}\tag{1.1}$$

(where the subscripts denote partial derivatives). They let  $z = x + iy$ ,  $f(z) = u + iv$  and introduced the  $\Sigma$ -derivative

$$\frac{d_{\Sigma} f}{d_{\Sigma} z} = f^{[1]}(z) = \sigma_1 u_x + i \frac{v_x}{\sigma_2} = \tau_1 v_y - i \frac{u_y}{\tau_2}\tag{1.2}$$

and the  $\Sigma$ -integral

$$\int_{z_0}^z f(z) d_{\Sigma} z = \int_{z_0}^z \sigma_2 u dx - \tau_2 v dy + i \left( \int_{z_0}^z \frac{1}{\sigma_1} v dx + \frac{1}{\tau_1} u dy \right).\tag{1.3}$$

Functions  $f(z)$ , where  $u$  and  $v$  are of class  $C^2$  satisfying (1.1) were called sigma-monogenic ( $\Sigma$ -monogenic), where

$$\Sigma = \begin{pmatrix} \sigma_1 & \tau_1 \\ \sigma_2 & \tau_2 \end{pmatrix}.$$

Then  $f^{[1]}(z)$  and  $\int_{z_0}^z f(z) d_{\Sigma} z$  are  $\Sigma'$ -monogenic where

$$\Sigma' = \begin{pmatrix} \frac{1}{\sigma_2} & \tau_1 \\ \frac{1}{\sigma_1} & \tau_2 \end{pmatrix}.$$

Clearly  $\Sigma'' = \Sigma$ .



They then introduced the generalized powers  $a \cdot Z^{(n)}(z_0; z)$ , where  $a$  is a complex constant, by the equations ( $\sim$  denotes class  $\Sigma'$ ):

$$\begin{aligned} a \cdot Z^{(0)}(z_0; z) &= a \cdot \tilde{Z}^{(0)}(z_0; z) = a \\ a \cdot Z^{(1)}(z_0; z) &= \int_{z_0}^z a \, d_{\Sigma'} \xi \\ a \cdot \tilde{Z}^{(1)}(z_0; z) &= \int_{z_0}^z a \, d_{\Sigma} \xi \\ a \cdot Z^{(n)}(z_0; z) &= n \int_{z_0}^z a \cdot \tilde{Z}^{(n-1)}(z_0; \xi) d_{\Sigma'} \xi \\ a \cdot \tilde{Z}^{(n)}(z_0; z) &= n \int_{z_0}^z a \cdot Z^{(n-1)}(z_0; \xi) d_{\Sigma} \xi . \end{aligned} \tag{1.4}$$

Bers and Gelbart showed that if  $f(z)$  is  $\Sigma$ -monogenic in a circle of radius  $R$  about  $z_0$ , ( $\sigma_i, \tau_i$  positive analytic) then

$$f(z) = \sum_{n=0}^{\infty} a_n \cdot Z^{(n)}(z_0; z) \tag{1.5}$$

for  $|z - z_0|$  sufficiently small. The  $a_n$  are uniquely determined by a "Taylor formula" as

$$a_n = \frac{f^{[n]}(z_0)}{n!}$$

where  $f^{[n]}(z)$  is the  $n$ -th  $\Sigma$ -derivative of  $f(z)$ .

In later work [2],  $\sigma_i$  and  $\tau_i$  were assumed to be positive and of class  $C^2$  whereas in [1],  $\sigma_i$  and  $\tau_i$  were analytic functions of their respective real variables.





One of the main unsolved problems in the theory of  $\Sigma$ -monogenic functions is:

If  $f(z)$  is  $\Sigma$ -monogenic in  $\Gamma = \{z; |z-z_0| < R\}$ , is

$$f(z) = \sum_{n=0}^{\infty} \frac{f^{[n]}(z_0)}{n!} \cdot z^{(n)}(z_0; z) \quad \text{in } \Gamma ?$$

We have succeeded in giving an affirmative answer to this question in the special case where  $\sigma_1 = \sigma_2 = e^{-cx}$ ,  $\tau_1 = \tau_2 = 1$ .

One important procedure is to set up a correspondence between analytic functions and  $\Sigma$ -monogenic functions; that is, given any analytic function  $f(z)$ , we can find explicitly a  $\Sigma$ -monogenic function (for our special case) which has some of the properties of  $f(z)$ . This is done by solving Cauchy problems on a parallel to the  $y$ -axis.

We consider then the special case

$$\begin{aligned} e^{-cx} u_x &= v_y \\ e^{-cx} u_y &= -v_x \end{aligned} \tag{1.6}$$



and, for convenience, assume that  $c > 0$ .

Assuming that  $u, v$  are of class  $C^2$ , it easily follows that

$$\Delta u - cu_x = u_{xx} + u_{yy} - cu_x = 0$$

$$\text{and } \Delta v + cv_x = 0.$$

The former equation is that satisfied by the vorticity in plane Oseen flow [3].

It is easy to see that the general equation with constant coefficients

$$\Delta u + a u_x + b u_y + c u = 0$$

can be reduced to this form provided  $a^2 + b^2 - 4c \geq 0$ . This is the case when the solution of the Dirichlet problem for the equation is unique [Paraf's Theorem].

In considering the ~~two~~ problem mentioned above we shall assume  $z_0 = 0$ . This is done merely to simplify the work: the general case can be handled in the same manner. Also the general formulae can be easily obtained from ours by a simple transformation.

For, let  $z_0 = x_0 + i y_0$  and suppose we are interested in equations (1.6) in a neighbourhood of  $z_0$ . Let  $X = x - x_0$ ,  $Y = y - y_0$ ,  $U = e^{-cx_0} u$ ,  $V = v$ . Then

$$e^{-cX} U_X = V_Y$$

$$e^{-cX} U_Y = -V_X$$





and so we can consider the last equation in a neighbourhood of  $X = Y = 0$ .

We now give the solutions of some Cauchy problems.

**THEOREM 1.1:** Suppose  $f(z)$  is holomorphic in a circle with centre the origin and radius  $R$ . Let  $D$  be the square whose vertices are  $(-R, 0)$ ,  $(0, R)$ ,  $(R, 0)$ , and  $(0, -R)$ .

i) Then, inside  $D$ , the function

$$u(x, y) = f(y) + \frac{x}{\pi} \int_0^1 e^{cux} (1-u)^{1/2} u^{-1/2} du \int_0^\pi i \cos \vartheta f'(y+2ix[u(1-u)])^{1/2} \cos \vartheta d\vartheta$$

is the solution of  $\Delta u - cu_x = 0$  satisfying the conditions

$$[u = f(y), \frac{\partial u}{\partial x} = 0] \text{ on } x = 0.$$

$$\text{ii) } u(x, y) = \frac{x}{\pi} \int_0^1 e^{cux} du \int_0^\pi f(y + 2ix[u(1-u)])^{1/2} \cos \vartheta d\vartheta$$

is the solution in  $D$  of  $\Delta u - cu_x = 0$ , satisfying

$$[u = 0, \frac{\partial u}{\partial x} = f(y)] \text{ on } x = 0.$$

iii) Inside  $D$ , the function

$$v(x, y) = f(y) + \frac{x}{\pi} \int_0^1 e^{-cux} (1-u)^{1/2} u^{-1/2} du \int_0^\pi i \cos \vartheta f'(y+2ix[u(1-u)])^{1/2} \cos \vartheta d\vartheta$$

is the solution of  $\Delta v + cv_x = 0$  satisfying the conditions

$$[v = f(y), \frac{\partial v}{\partial x} = 0] \text{ on } x = 0.$$

$$\text{iv) } v(x, y) = \frac{x}{\pi} \int_0^1 e^{-cux} du \int_0^\pi f(y + 2ix[u(1-u)])^{1/2} \cos \vartheta d\vartheta$$

is the solution in  $D$  of  $\Delta v + cv_x = 0$ , satisfying

$$[v = 0, \frac{\partial v}{\partial x} = f(y)] \text{ on } x = 0.$$



PROOF: We shall prove i) as the proofs of iii), iv) are the same as those for i), ii). ii) is proved in a similar fashion to i) and so we only point out the main differences.

i) First, it is clear that the  $u$  defined in i) above satisfies the Cauchy data.

To prove that it satisfies the equation, note that

$$\begin{aligned} f'(y + 2ix[u(1-u)]^{1/2} \cos \varnothing) &= f'(y) + (2ix[u(1-u)]^{1/2} \cos \varnothing) f''(y) + \dots \\ &+ \frac{(2ix[u(1-u)]^{1/2} \cos \varnothing)^n}{n!} f^{(n+1)}(y) + \dots \end{aligned}$$

provided  $(x, y) \in D$ .

$$\text{So, since } \int_0^\pi \cos^{2n+1} \varnothing \, d\varnothing = 0, \quad \int_0^\pi \cos^{2n} \varnothing \, d\varnothing = \pi \frac{(2n)!}{2^{2n}(n!)^2},$$

we get

$$\begin{aligned} u &= f(y) + \int_0^1 e^{cux} \, du \left[ \sum_{n=1}^{\infty} (-1)^n (1-u)^n u^{n-1} \frac{x^{2n}}{n!(n-1)!} f^{(2n)}(y) \right] \\ &= \sum_{n=0}^{\infty} (-1)^n a_{2n}(x) f^{(2n)}(y) \end{aligned}$$

where  $a_0 = 1$ ,

$$a_{2n}(x) = \frac{x^{2n}}{n!(n-1)!} \int_0^1 e^{cux} (1-u)^n u^{n-1} \, du.$$

The term-by-term integration in the above is valid in  $D$  as can be seen by using Cauchy's inequality for  $f^{(2n)}(y)$ .



Note also that

$$|a_{2n}(x)| \leq \frac{|x|^{2n}}{(2n)!} e^{|cx|}$$

since  $\int_0^1 (1-u)^n u^{n-1} du = \frac{n!(n-1)!}{(2n)!} .$

Now, if  $D \equiv \frac{d}{dx} ,$

$$(D^2 - cD)a_2 = 1 , \quad (D^2 - cD)a_{2n+2} = a_{2n} .$$

To show this write

$$a_{2n}(x) = \frac{1}{n!(n-1)!} \int_0^x (x-t)^n t^{n-1} e^{ct} dt$$

and do an integration by parts after differentiating.

Finally then, differentiating

$$u = \sum_{n=0}^{\infty} (-1)^n a_{2n}(x) f^{(2n)}(y)$$

term-by-term it is easily seen that  $\Delta u - cu_x = 0 .$

ii) The proof of ii) proceeds as above. The only difference is that instead of  $a_{2n}$  we have

$$\begin{aligned} A_0 &= x \int_0^1 e^{cux} du = \int_0^x e^{ct} dt , \\ A_{2n} &= \frac{x^{2n+1}}{(n!)^2} \int_0^1 e^{cux} u^n (1-u)^n du \\ &= \frac{1}{(n!)^2} \int_0^x e^{ct} t^n (x-t)^n dt . \end{aligned}$$

Again,  $(D^2 - cD)A_{2n} = A_{2n-2} , \quad (n \geq 1) .$

Q.E.D.





COROLLARY 1.1: If  $f(z)$  is an entire function, then the  $u$ 's and  $v$ 's defined above are solutions in the whole plane. If  $f(z)$  has a singularity at  $z = 0$ , then  $u$  and  $v$  are solutions for  $|x| < |y|$ .

We are now in a position to express our  $\Sigma$ -monogenic functions in terms of an analytic function.

THEOREM 1.2: i) Suppose  $f(z)$  is holomorphic in a circle  $|z| < R$ . Then, in the square  $D$  defined in Theorem 1.1,

$$u_1 + iv_1 = f(y) + \frac{x}{\pi} \int_0^1 e^{cux} (1-u)^{1/2} u^{-1/2} du \int_0^\pi i \cos \vartheta f'(y+2ix[u(1-u)])^{1/2} \cos \vartheta d\vartheta \\ - i \frac{x}{\pi} \int_0^1 e^{-cux} du \int_0^\pi f'(y+2ix[u(1-u)])^{1/2} \cos \vartheta d\vartheta$$

is  $\Sigma$ -monogenic and satisfies the Cauchy conditions

$$u_1(x=0) = f(y), \quad \frac{\partial v_1(x=0)}{\partial x} = -f'(y), \quad \frac{\partial u_1(x=0)}{\partial x} = v_1(x=0) = 0.$$

ii) Under the above assumptions about domains,

$$u_2 + iv_2 = \frac{x}{\pi} \int_0^1 e^{cux} du \int_0^\pi g'(y+2ix[u(1-u)])^{1/2} \cos \vartheta d\vartheta \\ + i \left\{ g(y) + \frac{x}{\pi} \int_0^1 e^{-cux} (1-u)^{1/2} u^{-1/2} du \int_0^\pi i \cos \vartheta g'(y+2ix \cdot \right. \\ \left. \cdot [u(1-u)]^{1/2} \cos \vartheta d\vartheta \right\}$$

is  $\Sigma$ -monogenic and satisfies the Cauchy data

$$u_2(x=0) = \frac{\partial v_2(x=0)}{\partial x} = 0, \quad v_2(x=0) = g(y), \\ \frac{\partial u_2(x=0)}{\partial x} = g'(y).$$



PROOF: i) As in Theorem 1.1, inside  $D$  we have

$$u = \sum_{n=0}^{\infty} (-1)^n a_{2n}(x) f^{(2n)}(y)$$

$$v = \sum_{n=0}^{\infty} (-1)^{n+1} \bar{A}_{2n}(x) f^{(2n+1)}(y) ,$$

where

$$a_0(x) = 1 ,$$

$$a_{2n}(x) = \frac{1}{n!(n-1)!} \int_0^x e^{ct} (x-t)^n t^{n-1} dt$$

and

$$\bar{A}_0 = \int_0^x e^{-ct} dt$$

$$\bar{A}_{2n-2} = \frac{1}{[(n-1)!]^2} \int_0^x e^{-ct} (x-t)^{n-1} t^{n-1} dt .$$

Now, we easily see that

$$\bar{A}_{2n-2} - e^{-cx} a'_{2n} = 0 \quad (n \geq 1) \quad \text{where ' denotes}$$

differentiation with respect to  $x$ .

So,

$$e^{-cx} u_x = v_y .$$

Also 
$$\bar{A}'_{2n} - e^{-cx} a_{2n} = 0 \quad (n \geq 0)$$

and thus

$$e^{-cx} u_y = -v_x .$$

ii) With  $u$  and  $v$  defined in ii) we have



$$u = \sum_{n=0}^{\infty} (-1)^n A_{2n}(x) g^{(2n+1)}(y)$$

$$v = \sum_{n=0}^{\infty} (-1)^n \bar{a}_{2n}(x) g^{(2n)}(y)$$

where

$$\bar{a}_0 = 1$$

$$\bar{a}_{2n} = \frac{1}{n!(n-1)!} \int_0^x e^{-ct} (x-t)^n t^{n-1} dt$$

and

$$A_0 = \int_0^x e^{ct} dt$$

$$A_{2n} = \frac{1}{(n!)^2} \int_0^x e^{ct} (x-t)^n t^n dt .$$

$$\text{Now, } e^{-cx} A'_{2n} - \bar{a}_{2n} = 0 \quad (n \geq 0) ,$$

$$\text{and so } e^{-cx} u_x = v_y .$$

$$\text{Also, } e^{-cx} A_{2n-2} - \bar{a}'_{2n} = 0 \quad (n \geq 1)$$

$$\text{and so } e^{-cx} u_y = -v_x . \quad \text{Q.E.D.}$$

We can now define a  $\Sigma$ -monogenic function "corresponding" to an analytic function.

DEFINITION 1.1: If  $F(z)$  is an analytic function in  $|z| < R$ , then the  $\Sigma$ -monogenic function corresponding to  $F(z)$  at  $z_0=0$  is

$$U + iV = (u_1 + iv_1) + (u_2 + iv_2)$$

where  $(u_1 + iv_1), (u_2 + iv_2)$  are defined in Theorem 1.2, and where





$$f(y) = \operatorname{Re} F(0 + iy)$$

$$g(y) = \operatorname{Im} F(0 + iy) .$$

$U + iV$  is defined in the square  $D$  described in Theorem 1.1.

For example, let  $F(z) = x + iy$ . Then  $f(y) = 0$ ,  $g(y) = y$ .

So

$$U + iV = 0 + \frac{x}{\pi} \int_0^1 e^{cux} du \int_0^\pi d\phi + i \left[ y + \frac{x}{\pi} \int_0^1 e^{-cux} (1-u)^{1/2} u^{-1/2} \cdot du \int_0^\pi i \cos \phi d\phi \right]$$

$$= \frac{1}{c} (e^{cx} - 1) + iy .$$

This is  $1 \cdot Z^{(1)}(0; z) = Z^{(1)}(0; z)$ .

Again, let  $F(z) = iz = ix - y$ . Then  $f(y) = -y$ ,  $g(y) = 0$ ,

and

$$U + iV = -y - \frac{x}{\pi} \int_0^1 e^{cux} (1-u)^{1/2} u^{-1/2} du \int_0^\pi i \cos \phi d\phi + \frac{x}{\pi} \int_0^1 e^{-cux} du \int_0^\pi d\phi$$

$$= -y + \frac{i}{c} (1 - e^{-cx}) .$$

This is  $i \cdot Z^{(1)}(0; z)$ .

We shall show that these examples indicate the general situation, namely that the  $\Sigma$ -monogenic function corresponding to  $z^n$  is  $Z^{(n)}(0; z)$  and the function corresponding to  $iz^n$  is  $i \cdot Z^{(n)}(0; z)$ . Thus the function corresponding to  $az^n$  is  $a \cdot Z^{(n)}(0; z)$ .



It therefore follows that if  $F(z) = \sum_{n=0}^{\infty} a_n z^n$ ,  $|z| < R$ ,

then the corresponding function is  $\sum_{n=0}^{\infty} a_n \cdot z^{(n)}(0; z)$ ,  $z \in D$ . Thus our definition of a corresponding function coincides with that of Bers and Gelbart.

Unfortunately, given a  $\Sigma$ -monogenic function  $U + iV$ , we do not know how to find the analytic function to which it corresponds. Thus we can not prove the Generalized Taylor's Theorem in this manner.



## CHAPTER II

### THE POWERS $a \cdot Z^{(n)}(z_0; z)$ AND THEIR BOUNDS.

We first give explicit formulae for the powers  $Z^{(n)}(z_0; z)$  and  $i \cdot Z^{(n)}(z_0; z)$  defined originally by Bers and Gelbart by equations (1.4). Then using these we show that

$$|a \cdot Z^{(n)}(0; z)| < 2^{3/2} (n+1) |a| e^{|cx|} r^n, \quad (r = (x^2 + y^2)^{1/2} = |z|).$$

We next show that if  $f(z)$  is  $\Sigma$ -monogenic in a circle  $|z| < R$ , where

$$\Sigma = \begin{pmatrix} e^{-cx} & 1 \\ e^{-cx} & 1 \end{pmatrix},$$

then

$$|f^{[n]}(0)| < \frac{K n!}{r^n}$$

where  $K$  does not depend on  $n$  and  $r < R$ . These bounds then suffice to show that  $\sum_{n=0}^{\infty} \frac{f^{[n]}(0)}{n!} \cdot Z^{(n)}(0; z)$  converges for  $|z| < R$  and it then follows (from the work of Bers and Gelbart) that  $f(z) = \sum_{n=0}^{\infty} \frac{f^{[n]}(0)}{n!} \cdot Z^{(n)}(0; z)$  for  $|z| < R$ .

As the first step in this programme we have the following result:

**THEOREM 2.1:** Suppose  $Z^{(n)}(z_0; z) = u^{(n)} + i v^{(n)}$   
and  $i \cdot Z^{(n)}(z_0; z) = \bar{u}^{(n)} + i \bar{v}^{(n)}$

where  $u^{(n)}, v^{(n)}, \bar{u}^{(n)}, \bar{v}^{(n)}$  are real and  $n$  is an integer  $\geq 0$ .

Then





$$u^{(2n)}(z_0; z) = (-1)^n (y-y_0)^{2n} + \frac{(-1)^n 2n(x-x_0)}{\pi} \int_0^1 e^{cu(x-x_0)} (1-u)^{1/2} u^{-1/2} \cdot$$

$$\left\{ \int_0^\pi i \cos \theta \left[ y-y_0 + 2i(x-x_0)[u(1-u)]^{1/2} \cos \theta \right]^{2n-1} d\theta \right\} du$$

$$= (-1)^n (y-y_0)^{2n} + (-1)^n 2(x-x_0)^2 \int_0^1 e^{cu(x-x_0)} (1-u) [(y-y_0)^2 +$$

$$4(x-x_0)^2 u(1-u)]^{n-1} \left\{ 2n P_{2n} \left( \frac{y-y_0}{[(y-y_0)^2 + 4(x-x_0)^2 u(1-u)]^{1/2}} \right) \right.$$

$$\left. - \frac{y-y_0}{[(y-y_0)^2 + 4(x-x_0)^2 u(1-u)]^{1/2}} P'_{2n} \left( \frac{y-y_0}{[(y-y_0)^2 + 4(x-x_0)^2 u(1-u)]^{1/2}} \right) \right\} du$$

where  $P_n(z)$  is Legendre's Polynomial.

$$u^{(2n+1)}(z_0; z) = \frac{(-1)^n e^{cx_0} (2n+1)(x-x_0)}{\pi} \int_0^1 e^{cu(x-x_0)} \left[ \int_0^\pi (y-y_0 + 2i(x-x_0)[u(1-u)]^{1/2} \cdot \right. \\ \left. \cdot \cos \theta \right)^{2n} d\theta \Big] du$$

$$= (-1)^n e^{cx_0} (2n+1)(x-x_0) \int_0^1 e^{cu(x-x_0)} [(y-y_0)^2 + 4(x-x_0)^2 u(1-u)]^n \cdot$$

$$\cdot P_{2n} \left( \frac{y-y_0}{[(y-y_0)^2 + 4(x-x_0)^2 u(1-u)]^{1/2}} \right) du :$$

$$v^{(2n)}(z_0; z) = e^{-cx_0} \frac{2n(-1)^{n-1} (x-x_0)}{\pi} \int_0^1 e^{-cu(x-x_0)} \left[ \int_0^\pi (y-y_0 + 2i(x-x_0) \cdot \right. \\ \left. [u(1-u)]^{1/2} \cos \theta \right)^{2n-1} d\theta \Big] du$$



$$= e^{-cx_0} (-1)^{n-1} 2n(x-x_0) \int_0^1 e^{-cu(x-x_0)} [(y-y_0)^2 + 4(x-x_0)^2 u(1-u)]^{n-1/2} \cdot P_{2n-1} \left( \frac{y-y_0}{[(y-y_0)^2 + 4(x-x_0)^2 u(1-u)]^{1/2}} \right) du .$$

$$v^{(2n+1)}(z_0; z) = (-1)^n (y-y_0)^{2n+1} + \frac{(-1)^n (x-x_0)(2n+1)}{\pi} \int_0^1 (1-u)^{1/2} u^{-1/2} \cdot$$

$$e^{-cu(x-x_0)} \left[ \int_0^\pi i \cos \theta [(y-y_0)+2i(x-x_0)[u(1-u)]^{1/2} \cos \theta]^{2n} d\theta \right] du$$

$$= (-1)^n (y-y_0)^{2n+1} + (-1)^n 2(x-x_0)^2 \int_0^1 e^{-cu(x-x_0)} (1-u) \cdot$$

$$[(y-y_0)^2 + 4(x-x_0)^2 u(1-u)]^{n-1/2} \left\{ (2n+1) \cdot$$

$$\cdot P_{2n+1} \left( \frac{y-y_0}{[(y-y_0)^2 + 4(x-x_0)^2 u(1-u)]^{1/2}} \right) - \frac{y-y_0}{[(y-y_0)^2 + 4(x-x_0)^2 u(1-u)]^{1/2}} \cdot$$

$$\cdot P'_{2n+1} \left( \frac{y-y_0}{[(y-y_0)^2 + 4(x-x_0)^2 u(1-u)]^{1/2}} \right) \Big\} du$$

The expressions for  $\frac{1}{u}^{(2n)}(z_0; z)$ ,  $\frac{1}{u}^{(2n+1)}(z_0; z)$ ,  $\frac{1}{v}^{(2n)}(z_0; z)$ ,  $\frac{1}{v}^{(2n+1)}(z_0; z)$  become obvious from Theorem 1.2.

PROOF: We note that the second form involving  $P_n(z)$  is easily obtained from the first by using Laplace's Integral

$$\frac{1}{\pi} \int_0^\pi [y-y_0 + i \zeta \cos \phi]^n d\phi = [(y-y_0)^2 + \zeta^2]^{n/2} P_n \left( \frac{y-y_0}{[(y-y_0)^2 + \zeta^2]^{1/2}} \right) .$$



The first forms for  $u^{(n)}$ ,  $v^{(n)}$  (and thus  $\bar{u}^{(n)}$ ,  $\bar{v}^{(n)}$ ) are obtained from Definition 1.1 and Theorem 1.2 by assuming that  $Z^{(n)}$  is the function corresponding to  $z^n$  and that  $i \cdot Z^{(n)}$  is the function corresponding to  $iz^n$ .

Thus Theorem 1.2 shows that these functions are  $\Sigma$ -monogenic in the whole plane.

Next, it is easily seen that for these functions

$$\frac{d_{\Sigma}}{d_{\Sigma} z} Z^{(n)}(z_0; z) = n \bar{Z}^{(n-1)}(z_0; z)$$

$$\frac{d_{\Sigma'}}{d_{\Sigma'} z} \bar{Z}^{(n)}(z_0; z) = n Z^{(n-1)}(z_0; z) \quad , \quad n \geq 1 .$$

Consider, for instance, the case  $n = 2p$ ,  $n$  an even integer.

$$\begin{aligned} \frac{d_{\Sigma}}{d_{\Sigma} z} Z^{(2p)}(z_0; z) &= \frac{\partial v^{(2p)}}{\partial y} - i \frac{\partial u^{(2p)}}{\partial y} \quad \text{from (1.2)} \\ &= \frac{2p(2p-1)}{\pi} \frac{e^{-cx_0} (-1)^{p-1} (x-x_0)}{\int_0^1 e^{-cu(x-x_0)} \cdot} \\ &\quad \left[ \int_0^{\pi} (y-y_0 + 2i(x-x_0)[u(1-u)])^{1/2} \cos \theta)^{2p-1} d\theta \right] du \\ &\quad - i \left\{ (-1)^p 2p(y-y_0)^{2p-1} + \frac{(-1)^p 2p(2p-1)(x-x_0)}{\pi} \right. \\ &\quad \left. \int_0^1 e^{cu(x-x_0)} (1-u)^{1/2} u^{-1/2} \left\{ \int_0^{\pi} i \cos \theta \cdot \right. \right. \\ &\quad \left. \left. \cdot [y-y_0 + 2i(x-x_0)[u(1-u)])^{1/2} \cos \theta \right]^{2p-2} d\theta \right\} du \Big\} \\ &= 2p \bar{Z}^{(2p-1)}(z_0; z) \end{aligned}$$





since, clearly,  $\tilde{Z}^{(n)}$  differs from  $Z^{(n)}$  only by changing  $c$  to  $-c$ .

Next, it is obvious that

$$\begin{aligned} Z^{(n)}(z_0; z_0) &= \tilde{Z}^{(n)}(z_0; z_0) = 0, \quad n \geq 1 \\ i \cdot Z^{(n)}(z_0; z_0) &= i \cdot \tilde{Z}^{(n)}(z_0; z_0) = 0, \quad n \geq 1. \end{aligned}$$

Thus, for  $n \geq 1$ ,

$$\begin{aligned} Z^{(n)}(z_0; z) &= n \int_{z_0}^z \tilde{Z}^{(n-1)}(z_0; \zeta) d_{\Sigma} \zeta \\ i \cdot Z^{(n)}(z_0; z) &= n \int_{z_0}^z i \cdot \tilde{Z}^{(n-1)}(z_0; \zeta) d_{\Sigma} \zeta \\ \tilde{Z}^{(n)}(z_0; z) &= n \int_{z_0}^z Z^{(n-1)}(z_0; \zeta) d_{\Sigma} \zeta \\ i \cdot \tilde{Z}^{(n)}(z_0; z) &= n \int_{z_0}^z i \cdot Z^{(n-1)}(z_0; \zeta) d_{\Sigma} \zeta. \end{aligned}$$

Also,

$$\begin{aligned} Z^{(0)}(z_0; z) &= \tilde{Z}^{(0)}(z_0; z) = 1 \\ i \cdot Z^{(0)}(z_0; z) &= i \cdot \tilde{Z}^{(0)}(z_0; z) = i. \end{aligned}$$

Thus the formulae we have given above for  $Z^{(n)}(z_0; z)$  and  $i \cdot Z^{(n)}(z_0; z)$  agree with those obtained by successive integration as in equations (1.4). Q.E.D.

COROLLARY 2.1: We can also define  $L(z_0; z)$  and  $i \cdot L(z_0; z)$  as the functions corresponding to  $\log(z - z_0)$  and  $i \log(z - z_0)$  respectively and  $Z^{(\alpha)}(z_0; z)$ ,  $i \cdot Z^{(\alpha)}(z_0; z)$  as the functions corresponding to  $(z - z_0)^\alpha$ ,  $i(z - z_0)^\alpha$  for  $\alpha$  any real number. These definitions give  $\Sigma$ -monogenic functions for  $y > y_0$  (or  $y < y_0$ ).



Then in  $y > y_0$  (or  $y < y_0$ )

$$\frac{d_{\Sigma} L(z_0; z)}{d_{\Sigma} z} = \gamma^{(-1)}(z_0; z)$$

$$\frac{d_{\Sigma} i \cdot L(z_0; z)}{d_{\Sigma} z} = i \cdot \gamma^{(-1)}(z_0; z)$$

$$\frac{d_{\Sigma} Z^{(\alpha)}(z_0; z)}{d_{\Sigma} z} = \alpha \gamma^{(\alpha-1)}(z_0; z)$$

$$\frac{d_{\Sigma} i \cdot Z^{(\alpha)}(z_0; z)}{d_{\Sigma} z} = \alpha i \cdot \gamma^{(\alpha-1)}(z_0; z) .$$

The proof is the same as in the above theorem.

Next, bounds for  $|Z^{(n)}(z_0; z)|$  and  $|i \cdot Z^{(n)}(z_0; z)|$  are obtained.

Without loss of generality, let  $z_0 = 0$ .

LEMMA 2.1: For  $z_0 = 0$ , and  $n$  an integer  $\geq 0$ ,

$$|u^{(2n)}| \leq e^{|cx|} (2n+1) r^{2n}$$

$$|u^{(2n+1)}| \leq e^{|cx|} (2n+1) r^{2n+1}$$

$$|v^{(2n)}| \leq e^{|cx|} (2n) r^{2n}$$

$$|v^{(2n+1)}| \leq e^{|cx|} (2n+2) r^{2n+1}$$

$$|\bar{u}^{(2n)}| \leq e^{|cx|} (2n) r^{2n}$$

$$|\bar{u}^{(2n+1)}| \leq e^{|cx|} (2n+2) r^{2n+1}$$

$$|\bar{v}^{(2n)}| \leq e^{|cx|} (2n+1) r^{2n}$$

$$|\bar{v}^{(2n+1)}| \leq e^{|cx|} (2n+1) r^{2n+1}$$



where  $r = (x^2 + y^2)^{1/2} = |z|$ .

PROOF: We will show the calculations involved in obtaining the bounds on  $u^{(2n)}(o; z)$ , as the others are analogous.

$$\begin{aligned} \text{Since } |y + 2ix[u(1-u)]^{1/2} \cos \theta|^2 &= y^2 + 4x^2 u(1-u) \cos^2 \theta \\ &\leq y^2 + x^2 \quad (0 \leq u \leq 1) \\ &= |x + iy|^2, \end{aligned}$$

$$\begin{aligned} \text{then } \left| \frac{1}{\pi} \int_0^\pi i \cos \theta (y + 2ix[u(1-u)]^{1/2} \cos \theta)^{2n-1} d\theta \right| &\leq \frac{1}{\pi} |x + iy|^{2n-1} \int_0^\pi |\cos \theta| d\theta \\ &= \frac{2}{\pi} |x + iy|^{2n-1}. \end{aligned}$$

Now from the first formula of Theorem 2.1 we have

$$\begin{aligned} u^{(2n)}(o; z) &= (-1)^n \left\{ y^{2n} + \frac{2nx}{\pi} \int_0^1 e^{cux} (1-u)^{1/2} u^{-1/2} du \int_0^\pi i \cos \theta \right. \\ &\quad \left. \cdot (y + 2ix [u(1-u)]^{1/2} \cos \theta)^{2n-1} d\theta \right\}. \end{aligned}$$

Therefore,

$$\begin{aligned} |u^{(2n)}(o; z)| &\leq y^{2n} + \frac{4n|x|}{\pi} e^{|cx|} |x + iy|^{2n-1} \int_0^1 (1-u)^{1/2} u^{-1/2} du \\ &= y^{2n} + 2n|x| e^{|cx|} |x + iy|^{2n-1} \\ &\leq e^{|cx|} |x + iy|^{2n-1} [|y| + 2n|x|]. \end{aligned}$$

Clearly,  $\max \{|y| + 2n|x|\}$  is  $r(4n^2 + 1)^{1/2}$ .

$$\begin{aligned} \text{Hence } |u^{(2n)}| &\leq e^{|cx|} |x + iy|^{2n} (4n^2 + 1)^{1/2} \\ &\leq e^{|cx|} (2n+1) r^{2n}. \end{aligned} \quad \text{Q.E.D.}$$





From this Lemma, the following theorem then follows.

THEOREM 2.2:  $|a \cdot Z^{(n)}(o; z)| < 2^{3/2(n+1)} |a| e^{|cx|} r^n$ ,  $n \geq 1$ ,

where  $r = (x^2 + y^2)^{1/2}$ .

PROOF: Let  $a = \alpha + i\beta$  where  $\alpha$  and  $\beta$  are real.

$$\begin{aligned} \text{Then, } |a \cdot Z^{(2n)}(o; z)| &= |\alpha Z^{(2n)}(o; z) + \beta i \cdot Z^{(2n)}(o; z)| \\ &= |\alpha u^{(2n)} + i \beta \bar{v}^{(2n)} + i \alpha v^{(2n)} + \beta \bar{u}^{(2n)}| \\ &\leq 2^{1/2} |a| [ |u^{(2n)}|^2 + |\bar{u}^{(2n)}|^2 + |v^{(2n)}|^2 + |\bar{v}^{(2n)}|^2 ]^{1/2} \\ &\quad \text{(by Schwarz's inequality)} \end{aligned}$$

$$\begin{aligned} &\leq 2 |a| e^{|cx|} r^{2n} \{ (2n+1)^2 + (2n)^2 \}^{1/2} \\ &< 2^{3/2} (2n+1) |a| e^{|cx|} r^{2n}. \end{aligned}$$

$$\begin{aligned} \text{Also, } |a \cdot Z^{(2n+1)}(o; z)| &= |\alpha u^{(2n+1)} + i \beta \bar{v}^{(2n+1)} + i \alpha v^{(2n+1)} + \beta \bar{u}^{(2n+1)}| \\ &\leq 2^{1/2} |a| [ |u^{(2n+1)}|^2 + |\bar{v}^{(2n+1)}|^2 + |v^{(2n+1)}|^2 + \\ &\quad + |\bar{u}^{(2n+1)}|^2 ]^{1/2} \\ &\leq 2^{1/2} |a| \{ 2(2n+1)^2 + 2(2n+2)^2 \}^{1/2} e^{|cx|} r^{2n+1} \\ &< 2^{3/2} |a| (2n+2) e^{|cx|} r^{2n+1}. \end{aligned}$$

Q.E.D.

COROLLARY 2.2:  $|a \cdot Z^{(n)}(z_o; z)| < 2^{3/2(n+1)} |a| e^{|cx| + |cx_o|} r^n$

where  $r = [(x-x_o)^2 + (y-y_o)^2]^{1/2}$  and  $n \geq 0$ .



PROOF: Everything is as with  $a \cdot Z^{(n)}(0; z)$  except that  $x$  is replaced by  $(x-x_0)$ ,  $y$  by  $(y-y_0)$  and there is a factor  $e^{cx_0}$  or  $e^{-cx_0}$ . Now,

$$\begin{aligned} |e^{cx_0 - cux_0}| &\leq e^{|cx_0|(1-u)}, & 0 \leq u \leq 1, \\ &\leq e^{|cx_0|} \end{aligned}$$

and so the result follows.



# CHAPTER III

## ESTIMATES ON THE DERIVATIVES OF $f(z)$ AND THE GENERALIZED TAYLOR'S THEOREM.

We wish to show that if  $f(z)$  is  $\Sigma$ -monogenic in  $|z| < R$ , then

$$f(z) = \sum_{n=0}^{\infty} a_n \cdot z^{(n)}(0; z)$$

where  $a_n = \frac{f^{[n]}(0)}{n!}$ .

Now we already have bounds on  $|z^{(n)}(0; z)|$  and so we shall now obtain bounds on  $f^{[n]}(0)$ .

Here,  $f^{[0]}(z) = f(z) = u + iv$

$$f^{[1]}(z) = \frac{d_{\Sigma} f}{d_{\Sigma} z} = v_y - i u_y$$

$$f^{[n]}(z) = \begin{cases} \frac{d_{\Sigma} f^{[n-1]}(z)}{d_{\Sigma} z}, & \text{if } n \text{ is odd} \\ \frac{d_{\Sigma'} f^{[n-1]}(z)}{d_{\Sigma'} z}, & \text{if } n \text{ is even.} \end{cases}$$

Hence,

$$f^{[2n]}(z) = (-1)^n \frac{\partial^{2n} u}{\partial y^{2n}} + i(-1)^n \frac{\partial^{2n} v}{\partial y^{2n}} \quad (3.1)$$

$$f^{[2n-1]}(z) = (-1)^{n-1} \frac{\partial^{2n-1} v}{\partial y^{2n-1}} + i(-1)^n \frac{\partial^{2n-1} u}{\partial y^{2n-1}}.$$

( $n$  a positive integer)





$$\left. \begin{array}{l} \text{Let } X = e^{-\frac{c}{2}x} u, \quad Y = e^{\frac{c}{2}x} v. \\ \text{Then, } \Delta X - \frac{c^2}{4} X = 0, \quad \Delta Y - \frac{c^2}{4} Y = 0. \end{array} \right\} \quad (3.2)$$

Bounds on the derivatives of  $X$  and  $Y$  will be found from the explicit solution of the Dirichlet Problem for  $\Delta X - \frac{c^2}{4} X = 0$  in  $|z| < R$ .

First, it is easily seen that  $K_0(\frac{c}{2}r)$ ,  $I_n(\frac{c}{2}r) \cos n\theta$ ,  $I_n(\frac{c}{2}r) \sin n\theta$ , ( $n \geq 1$ ) are solutions of  $\Delta X - \frac{c^2}{4} X = 0$ , where  $(r, \theta)$  are polar coordinates in the  $z$ -plane with origin  $z = 0$ .

Therefore,  $K_0(\frac{c}{2}[r^2 + R^2 - 2rR \cos(\theta - \phi)]^{1/2})$  is also a solution.

Here,  $I_n(z)$ ,  $K_n(z)$  are Bessel functions of pure imaginary argument [4, pp. 77-78].

$$K_0(\frac{c}{2}r) = -\log(\frac{c}{4}r) - \gamma + O(r \log r), \quad \text{for } r \text{ small.}$$

Using these solutions we can obtain a Green's function.

LEMMA 3.1: Let  $G(\text{Re}^{i\phi}, \text{re}^{i\theta})$  be the Green's function for  $\Delta u - \frac{c^2}{4} u = 0$ ; that is a solution of the equation with respect to  $r$  and  $\theta$  in  $|z| < a$ , which is 0 on  $|z| = a$  and which has a singularity  $-\log |PP'|$  for  $P = \text{re}^{i\theta}$  near  $P' = \text{Re}^{i\phi}$ . Then



$$\begin{aligned}
 G(Re^{i\phi}, re^{i\theta}) &= K_0\left(\frac{c}{2}[r^2 + R^2 - 2Rr \cos(\theta-\phi)]^{1/2}\right) \\
 &- \frac{K_0\left(\frac{c}{2}a\right)}{I_0\left(\frac{c}{2}a\right)} I_0\left(\frac{c}{2}r\right) I_0\left(\frac{c}{2}R\right) \\
 &- 2 \sum_{n=1}^{\infty} \frac{I_n\left(\frac{c}{2}r\right)}{I_n\left(\frac{c}{2}a\right)} K_n\left(\frac{c}{2}a\right) I_n\left(\frac{c}{2}R\right) \cos n(\theta-\phi)
 \end{aligned} \tag{3.3}$$

where  $R < a$ .

PROOF: From the asymptotic properties of  $I_n$  and  $K_n$  it is clear that the above series converges for  $R < a$ , and it can be differentiated term by term any number of times.

So the above  $G$  is a solution of the equation and it obviously has a singularity  $-\log |PP'|$ .

Next,

$$\begin{aligned}
 G(Re^{i\phi}, ae^{i\theta}) &= K_0\left(\frac{c}{2}[a^2 + R^2 - 2aR \cos(\theta-\phi)]^{1/2}\right) \\
 &- K_0\left(\frac{c}{2}a\right) I_0\left(\frac{c}{2}R\right) \\
 &- 2 \sum_{n=1}^{\infty} K_n\left(\frac{c}{2}a\right) I_n\left(\frac{c}{2}R\right) \cos n(\theta-\phi).
 \end{aligned}$$

But this is zero by [4, p. 361].

Q.E.D.

LEMMA 3.2: If  $\Delta X - \frac{c^2}{4} X = 0$  for  $R < a$  where  $x = R \cos \phi$ ,  $y = R \sin \phi$ , and if  $X$  is  $C^{(2)}$  for  $R \leq a$ , then



$$X(Re^{i\phi}) = \frac{1}{2\pi} \int_0^{2\pi} \left[ \frac{I_0(\frac{c}{2} R)}{I_0(\frac{c}{2} a)} + 2 \sum_{n=1}^{\infty} \frac{I_n(\frac{c}{2} R)}{I_n(\frac{c}{2} a)} \cos n(\theta - \phi) \right] f(\theta) d\theta \quad (3.4)$$

for  $R < a$ , where  $X(ae^{i\theta}) = f(\theta)$ .

PROOF: In the usual manner, we can express  $X(Re^{i\phi})$  by means of an integral over the boundary of  $|z| = a$ .

$$X(Re^{i\phi}) = -\frac{a}{2\pi} \int_0^{2\pi} X(ae^{i\theta}) \left[ \frac{\partial G(Re^{i\phi}, re^{i\theta})}{\partial r} \right]_{r=a} d\theta.$$

$$\text{Since } K'_0(x) = -K_1(x), \quad I'_0(x) = I_1(x),$$

$$-2K'_n(x) = K_{n-1}(x) + K_{n+1}(x),$$

$$2I'_n(x) = I_{n-1}(x) + I_{n+1}(x),$$

we have

$$\begin{aligned} \frac{\partial G(Re^{i\phi}, re^{i\theta})}{\partial r} &= -\frac{\frac{c}{2}(r-R \cos(\theta-\phi))}{[R^2+r^2-2rR \cos(\theta-\phi)]^{1/2}} K_1\left(\frac{c}{2}[R^2+r^2-2rR \cos(\theta-\phi)]^{1/2}\right) \\ &- \frac{c}{2} \frac{K_0(\frac{c}{2}a)}{I_0(\frac{c}{2}a)} I_1\left(\frac{c}{2}r\right) I_0\left(\frac{c}{2}R\right) \\ &- \frac{c}{2} \sum_{n=1}^{\infty} \frac{K_n(\frac{c}{2}a)}{I_n(\frac{c}{2}a)} I_n\left(\frac{c}{2}R\right) [I_{n-1}\left(\frac{c}{2}r\right) + I_{n+1}\left(\frac{c}{2}r\right)] \cos n(\theta-\phi). \end{aligned}$$

But, [4, p. 361]



$$\frac{a-R \cos (\theta-\phi)}{[a^2+R^2-2aR \cos (\theta-\phi)]^{1/2}} K_1\left(\frac{c}{2}\left[a^2+R^2-2aR \cos (\theta-\phi)\right]^{1/2}\right)$$

$$= K_1\left(\frac{c}{2} a\right) I_0\left(\frac{c}{2} R\right) + \sum_{1}^{\infty} I_n\left(\frac{c}{2} R\right)\left[K_{n+1}\left(\frac{c}{2} a\right)+K_{n-1}\left(\frac{c}{2} a\right)\right] \cos n(\theta-\phi)$$

and so

$$\left[\frac{\partial G\left(\operatorname{Re}^{i \phi}, \operatorname{re}^{i \theta}\right)}{\partial r}\right]_{r=a}=-\frac{c}{2} \frac{I_0\left(\frac{c}{2} R\right)}{I_0\left(\frac{c}{2} a\right)}\left[K_0\left(\frac{c}{2} a\right) I_1\left(\frac{c}{2} a\right)+K_1\left(\frac{c}{2} a\right) I_0\left(\frac{c}{2} a\right)\right]$$

$$-\frac{c}{2} \sum_{1}^{\infty} \frac{I_n\left(\frac{c}{2} R\right)}{I_n\left(\frac{c}{2} a\right)}\left[K_n\left(\frac{c}{2} a\right) I_{n-1}\left(\frac{c}{2} a\right)+K_n\left(\frac{c}{2} a\right) I_{n+1}\left(\frac{c}{2} a\right)\right.$$

$$\left.+\left.I_n\left(\frac{c}{2} a\right) K_{n+1}\left(\frac{c}{2} a\right)+I_n\left(\frac{c}{2} a\right) K_{n-1}\left(\frac{c}{2} a\right)\right] \cos n(\theta-\phi) .$$

However, [4, p. 80]

$$I_n(x) K_{n+1}(x)+I_{n+1}(x) K_n(x)=\frac{1}{x}$$

and so

$$\left[\frac{\partial G\left(\operatorname{Re}^{i \phi}, \operatorname{re}^{i \theta}\right)}{\partial r}\right]_{r=a}=-\frac{1}{a}\left\{\frac{I_0\left(\frac{c}{2} R\right)}{I_0\left(\frac{c}{2} a\right)}+2 \sum_{1}^{\infty} \frac{I_n\left(\frac{c}{2} R\right)}{I_n\left(\frac{c}{2} a\right)} \cos n(\theta-\phi)\right\} .$$

Substituting this expression into our initial expression for  $X\left(\operatorname{Re}^{i \phi}\right)$ , we get (3.4). Q.E.D.

We can obtain some interesting corollaries from (3.4) which we set down here even though they will not be used subsequently.





COROLLARY 3.1: The Mean Value Theorem.

$$X(R=0) = \frac{1}{2\pi I_0\left(\frac{c}{2}a\right)} \int_0^{2\pi} X(ae^{i\theta}) d\theta .$$

(Obvious).

COROLLARY 3.2: The Maximum Principle.

$$|X(R=0)| \leq \frac{\max_{\theta} |X(ae^{i\theta})|}{I_0\left(\frac{c}{2}a\right)} < \max_{\theta} |X(ae^{i\theta})| .$$

COROLLARY 3.3: If  $X$  is a solution in the whole plane of  $\Delta u - \frac{c^2}{4}u = 0$ ,

and

$$X = o\left(\frac{e^{\frac{c}{2}R}}{R^{1/2}}\right) \quad \text{for } R \text{ large,}$$

then  $X \equiv 0$ .

COROLLARY 3.4: There are no entire solutions bounded at infinity of  $\Delta u - \frac{c^2}{4}u = 0$  except  $u \equiv 0$ . In fact, there are no polynomial solutions except  $u \equiv 0$ .

Using equation (3.4) we can now obtain expressions for the derivatives of  $X$ . We first prove the following lemma.



LEMMA 3.3:

$$\left[ \frac{\partial^p (I_n(\frac{c}{2} r) \cos n\theta)}{\partial y^p} \right]_{x=y=0} = \begin{cases} \frac{(-1)^{\frac{n}{2}} p! (\frac{c}{4})^p}{(\frac{p-n}{2})! (\frac{p+n}{2})!} , & p \text{ even and } n \text{ even} \\ & \text{and } p \geq n > 0 . \\ 0 & , p \text{ odd or } n \text{ odd or} \\ & p < n . \end{cases}$$

$$\left[ \frac{\partial^p (I_n(\frac{c}{2} r) \sin n\theta)}{\partial y^p} \right]_{x=y=0} = \begin{cases} \frac{(-1)^{\frac{n-1}{2}} p! (\frac{c}{4})^p}{(\frac{p-n}{2})! (\frac{p+n}{2})!} , & p \text{ odd and } n \text{ odd and} \\ & p \geq n > 0 . \\ 0 & , p \text{ even or } n \text{ even or} \\ & p < n . \end{cases}$$

$$\left[ \frac{\partial^p I_0(\frac{c}{2} r)}{\partial y^p} \right]_{x=y=0} = \begin{cases} 0 & , p \text{ odd} \\ (\frac{c}{4})^p \frac{p!}{[(\frac{p}{2})!]^2} , & p \text{ even} , \end{cases}$$

where  $x = r \cos \theta$  ,  $y = r \sin \theta$  .

PROOF: Take the last relation first and use the result [4, pp. 360-361]

$$I_0(\frac{c}{2} r) = I_0(\frac{c}{2} x) I_0(\frac{c}{2} y) + 2 \sum_{m=1}^{\infty} (-1)^m I_m(\frac{c}{2} x) I_m(\frac{c}{2} y) \cos \frac{m\pi}{2} .$$

Then the result follows immediately since

$$I_m(x=0) = 0 , \quad (m \geq 1) \quad \text{and}$$

$$I_0(\frac{c}{2} y) = \sum_{m=0}^{\infty} \frac{(\frac{c}{4})^{2m} y^{2m}}{(m!)^2} .$$



The other results follow in the same manner using

$$I_n\left(\frac{c}{2} r\right) \cos n\theta = I_n\left(\frac{c}{2} x\right) I_0\left(\frac{c}{2} y\right) + \sum_{m=1}^{\infty} (-1)^m \cos \frac{m\pi}{2} I_m\left(\frac{c}{2} y\right) [I_{m+n}\left(\frac{c}{2} x\right) + I_{m-n}\left(\frac{c}{2} x\right)] .$$

$$I_n\left(\frac{c}{2} r\right) \sin n\theta = \sum_{m=1}^{\infty} (-1)^m \sin \frac{m\pi}{2} I_m\left(\frac{c}{2} y\right) [I_{m+n}\left(\frac{c}{2} x\right) - I_{m-n}\left(\frac{c}{2} x\right)] .$$

Q.E.D.

LEMMA 3.4: For  $r \geq 1$ ,

$$\left| \left[ \frac{\partial^{2r-1} X(Re^{i\theta})}{\partial y^{2r-1}} \right]_{R=0} \right| \leq \frac{4}{\pi} M \left(\frac{c}{4}\right)^{2r-1} (2r-1)!, \sum_{m=1}^r \frac{1}{(r-m)!(r+m-1)! I_{2m-1}\left(\frac{ca}{2}\right)}$$

$$\left| \left[ \frac{\partial^{2r} X(Re^{i\theta})}{\partial y^{2r}} \right]_{R=0} \right| \leq \frac{4}{\pi} M \left(\frac{c}{4}\right)^{2r} (2r)!, \sum_{m=0}^r \frac{1}{(r-m)!(r+m)! I_{2m}\left(\frac{ca}{2}\right)}$$

where  $M = \text{Maximum } |X(ae^{i\theta})|$ .

As before we are assuming that  $\Delta X - \frac{c^2}{4} X = 0$  in  $R < a$  and  $X \in C^{(2)}$  for  $R \leq a$ .

PROOF: i) Differentiating (3.4) and using Lemma 3.3 we get, for  $r \geq 1$ ,

$$\left[ \frac{\partial^{2r-1} X(Re^{i\theta})}{\partial y^{2r-1}} \right]_{R=0} = \frac{1}{\pi} \int_0^{2\pi} f(\theta) \left\{ \sum_{\substack{n=1 \\ (n \text{ odd})}}^{2r-1} \frac{(-1)^{\frac{n-1}{2}} (2r-1)! \left(\frac{c}{4}\right)^{2r-1}}{\left(\frac{2r-1-n}{2}\right)! \left(\frac{2r+n-1}{2}\right)! I_n\left(\frac{ca}{2}\right)} \sin n\theta \right\} d\theta$$





$$= \frac{1}{\pi} \int_0^{2\pi} f(\theta) \left\{ \sum_{m=1}^r \frac{(-1)^m (2r-1)! \left(\frac{c}{4}\right)^{2r-1}}{(r-m)! (r+m-1)! I_{2m-1}\left(\frac{ca}{2}\right)} \sin (2m-1)\theta \right\} d\theta$$

and so

$$\begin{aligned} \left| \left[ \frac{\partial^{2r-1} X(\text{Re } i\theta)}{\partial y^{2r-1}} \right]_{R=0} \right| &\leq \frac{(2r-1)! \left(\frac{c}{4}\right)^{2r-1}}{\pi} \sum_1^r \frac{\int_0^{2\pi} M |\sin(2m-1)\theta| d\theta}{(r-m)! (r+m-1)! I_{2m-1}\left(\frac{ca}{2}\right)} \\ &= \frac{4}{\pi} M (2r-1)! \left(\frac{c}{4}\right)^{2r-1} \sum_1^r \frac{1}{(r-m)! (r+m-1)! I_{2m-1}\left(\frac{ca}{2}\right)} \end{aligned}$$

where we have used  $\int_0^{2\pi} |\sin (2m-1)\theta| d\theta = 4$ .

Note that using Schwarz's inequality on  $\int_0^{2\pi} f(\theta) \sin (2m-1)\theta d\theta$  gives a worse result since  $2^{1/2} \pi > 4$ .

ii) Again, using equation (3.4) and Lemma 3.3, we get

$$\left| \left[ \frac{\partial^{2r} X(\text{Re } i\theta)}{\partial y^{2r}} \right]_{R=0} \right| = \left| \frac{1}{2\pi} \int_0^{2\pi} \left\{ \frac{\left(\frac{c}{4}\right)^{2r} (2r)!}{(r!)^2 I_0\left(\frac{ca}{2}\right)} + 2 \sum_{\substack{n=2 \\ (n \text{ even})}}^{2r} \frac{(-1)^{\frac{n}{2}} (2r)! \left(\frac{c}{4}\right)^{2r}}{\left(\frac{2r-n}{2}\right)! \left(\frac{2r+n}{2}\right)! I_n\left(\frac{ca}{2}\right)} \cos n\theta \right\} \cdot f(\theta) d\theta \right|$$

$$= \frac{\left(\frac{c}{4}\right)^{2r} (2r)!}{2\pi} \left| \int_0^{2\pi} \left\{ \frac{1}{(r!)^2 I_0\left(\frac{ca}{2}\right)} + \right. \right.$$

$$\left. 2 \sum_1^r \frac{(-1)^m \cos 2m\theta}{(r-m)! (r+m)! I_{2m}\left(\frac{ca}{2}\right)} \right\} f(\theta) d\theta \right|$$



$$\leq M\left(\frac{c}{4}\right)^{2r} (2r)! \left\{ \frac{1}{(r!)^2 I_0\left(\frac{ca}{2}\right)} + \frac{4}{\pi} \sum_{m=1}^r \frac{1}{(r-m)! (r+m)! I_{2m}\left(\frac{ca}{2}\right)} \right\}$$

$$< \frac{4}{\pi} M\left(\frac{c}{4}\right)^{2r} (2r)! \sum_{m=0}^r \frac{1}{(r-m)! (r+m)! I_{2m}\left(\frac{ca}{2}\right)} .$$

Q.E.D.

We next obtain bounds on the series

$$\sum_{m=1}^r \frac{1}{(r-m)! (m+r-1)! I_{2m-1}\left(\frac{ca}{2}\right)}$$

and

$$\sum_{m=0}^r \frac{1}{(r-m)! (r+m)! I_{2m}\left(\frac{ca}{2}\right)} .$$

LEMMA 3.5:

$$\left| \left[ \frac{\partial^{2r-1} X(\operatorname{Re} i\theta)}{\partial y^{2r-1}} \right]_{R=0} \right| \leq \frac{4}{\pi} \frac{M[(2r-1)!]}{a^{2r-1}} \left[ \frac{4}{ac} I_1\left(\frac{ac}{2}\right) \right] .$$

$$\left| \left[ \frac{\partial^{2r} X(\operatorname{Re} i\theta)}{\partial y^{2r}} \right]_{R=0} \right| \leq \frac{4}{\pi} \frac{M(2r)!}{a^{2r}} I_0\left(\frac{ac}{2}\right) .$$

PROOF: i) Consider first

$$\sum_{m=1}^r \frac{1}{(r-m)! (m+r-1)! I_{2m-1}\left(\frac{ca}{2}\right)}$$

$$= \frac{1}{(2r-1)! I_{2r-1}\left(\frac{ac}{2}\right)} \sum_{m=1}^r \frac{(2r-1)! I_{2r-1}\left(\frac{ac}{2}\right)}{(r-m)! (m+r-1)! I_{2m-1}\left(\frac{ac}{2}\right)} .$$



Now from [4, p. 153], or alternatively from the power series for  $I_n(x)$  for  $x$  real and  $n > 0$ ,

$$\frac{I_{n+2}(x)}{I_n(x)} = 1 - \frac{2(n+1)}{2(n+1) + \frac{x^2}{2(n+2) + \dots}} \leq \frac{x^2}{4(n+1)(n+2)}.$$

Therefore

$$\frac{I_{2r-1}(\frac{ac}{2})}{I_{2m-1}(\frac{ac}{2})} = \frac{I_{2r-1}(\frac{ac}{2})}{I_{2r-3}(\frac{ac}{2})} \cdot \frac{I_{2r-3}(\frac{ac}{2})}{I_{2r-5}(\frac{ac}{2})} \dots \frac{I_{2m+1}(\frac{ac}{2})}{I_{2m-1}(\frac{ac}{2})} \leq \left(\frac{ac}{4}\right)^{2r-2m} \frac{(2m-1)!}{(2r-1)!}.$$

$$\text{and so } \sum_1^r \frac{1}{(r-m)!(m+r-1)! I_{2m-1}(\frac{ac}{2})} \leq \frac{1}{(2r-1)! I_{2r-1}(\frac{ac}{2})} \sum_1^r \frac{(2m-1)!}{(r-m)!(m+r-1)!} \cdot \left(\frac{ac}{4}\right)^{2r-2m}$$

$$= \frac{1}{(2r-1)! I_{2r-1}(\frac{ac}{2})} \left\{ 1 + \frac{(\frac{ac}{4})^2}{1!(2r-2)} + \frac{(\frac{ac}{4})^4}{2!(2r-3)(2r-4)} + \dots + \frac{(\frac{ac}{4})^{2r-2}}{(r-1)! r!} \right\}$$

$$= \frac{1}{(2r-1)! I_{2r-1}(\frac{ac}{2})} \sum_{q=0}^{r-1} \frac{(\frac{ac}{4})^{2q}}{q! [(2r-q-1) \dots (2r-2q)]}.$$



Now in the general term of the above series,  $r-1 \geq q$  and so  $2r - 2q \geq 2$ .

$$\begin{aligned}
 \text{Therefore } \sum_{q=0}^{r-1} \frac{\left(\frac{ac}{4}\right)^{2q}}{q! [(2r-q-1) \dots (2r-2q)]} &\leq \sum_{q=0}^{r-1} \frac{\left(\frac{ac}{4}\right)^{2q}}{q! (q+1)!} \\
 &\leq \sum_{q=0}^{\infty} \frac{\left(\frac{ac}{4}\right)^{2q}}{q! (q+1)!} \\
 &= \frac{4}{ac} \sum_{q=0}^{\infty} \frac{\left(\frac{ac}{4}\right)^{2q+1}}{q! (q+1)!} \\
 &= \frac{4}{ac} I_1\left(\frac{ac}{2}\right) .
 \end{aligned}$$

Thus we have

$$\begin{aligned}
 \sum_{m=1}^r \frac{1}{(r-m)! (m+r-1)! I_{2m-1}\left(\frac{ac}{2}\right)} &\leq \frac{4}{ac} \frac{I_1\left(\frac{ac}{2}\right)}{(2r-1)! I_{2r-1}\left(\frac{ac}{2}\right)} \\
 &\leq \frac{I_1\left(\frac{ac}{2}\right)}{\left(\frac{ac}{4}\right)^{2r}}
 \end{aligned}$$

and so from Lemma 3.4,

$$\begin{aligned}
 \left| \left[ \frac{\partial^{2r-1} X(\text{Re } i\theta)}{\partial y^{2r-1}} \right]_{R=0} \right| &\leq \frac{4}{\pi} M(2r-1)! \left(\frac{c}{4}\right)^{2r-1} \frac{4 I_1\left(\frac{ac}{2}\right)}{ac (2r-1)! I_{2r-1}\left(\frac{ac}{2}\right)} \\
 &\leq \frac{4M}{\pi a} \frac{(2r-1)!}{2^{2r-1}} \frac{I_1\left(\frac{ac}{2}\right)}{\left(\frac{ac}{4}\right)} .
 \end{aligned}$$





ii) Now consider

$$\sum_0^r \frac{1}{(r-m)!(r+m)! I_{2m}(\frac{ca}{2})} = \frac{1}{(2r)! I_{2r}(\frac{ac}{2})} \sum_0^r \frac{(2r)! I_{2r}(\frac{ac}{2})}{(r-m)!(r+m)! I_{2m}(\frac{ca}{2})} .$$

Again, as before

$$\frac{(2r)! I_{2r}(\frac{ac}{2})}{I_{2m}(\frac{ac}{2})} \leq (\frac{ac}{4})^{2r-2m} (2m)!$$

and so

$$\begin{aligned} \sum_0^r \frac{1}{(r-m)!(r+m)! I_{2m}(\frac{ac}{2})} &\leq \frac{1}{(2r)! I_{2r}(\frac{ac}{2})} \sum_0^r \frac{(\frac{ac}{4})^{2r-2m} (2m)!}{(r-m)!(r+m)!} \\ &= \frac{1}{(2r)! I_{2r}(\frac{ac}{2})} \left\{ 1 + \frac{(\frac{ac}{4})^2}{1!(2r-1)} + \frac{(\frac{ac}{4})^4}{2!(2r-2)(2r-3)} \right. \\ &\quad \left. + \dots + \frac{(\frac{ac}{4})^{2r}}{r! r!} \right\} \\ &= \frac{1}{(2r)! I_{2r}(\frac{ac}{2})} \sum_0^r \frac{(\frac{ac}{4})^{2q}}{q! [(2r-q) \dots (2r-2q+1)]} \\ &\leq \frac{1}{(2r)! I_{2r}(\frac{ac}{2})} \sum_0^\infty \frac{(\frac{ac}{4})^{2q}}{q! q!} \\ &= \frac{1}{(2r)! I_{2r}(\frac{ac}{2})} I_0(\frac{ac}{2}) \\ &\leq \frac{I_0(\frac{ac}{2})}{(\frac{ac}{4})^{2r}} . \end{aligned}$$

Q.E.D.



We can now formulate the following theorem on  $f^{[n]}(0)$ .

**THEOREM 3.1:** If  $f(z)$  is  $\Sigma$ -monogenic in  $|z| \leq a$ , with

$$\Sigma = \begin{pmatrix} e^{-cx} & 1 \\ e^{-cx} & 1 \end{pmatrix},$$

then for  $n \geq 1$ ,

$$|f^{[2n]}(0)| \leq \frac{4}{\pi} \frac{2^{1/2} M(f)}{a^{2n}} (2n)! e^{\left|\frac{ca}{2}\right|} I_0\left(\frac{ac}{2}\right)$$

$$|f^{[2n-1]}(0)| \leq \frac{4}{\pi} \frac{2^{1/2} M(f)}{a^{2n-1}} (2n-1)! e^{\left|\frac{ca}{2}\right|} \frac{4}{ac} I_1\left(\frac{ac}{2}\right),$$

where  $M(f) = \text{Maximum } |f(z)|$   
( $|z|=a$ )

Alternatively,

$$|f^{[n]}(0)| \leq \frac{4}{\pi} \frac{M(f)}{a^n} 2^{1/2} n! e^{\left|\frac{ca}{2}\right|} I_0\left(\frac{ac}{2}\right). \quad (3.5)$$

**PROOF:** Using equations (3.1) and (3.2) and Lemma 3.5,

$$|f^{[2n]}(0)| \leq \left\{ \left( \left[ \frac{\partial^{2n} u}{\partial y^{2n}} \right]_{z=0} \right)^2 + \left( \left[ \frac{\partial^{2n} v}{\partial y^{2n}} \right]_{z=0} \right)^2 \right\}^{1/2}$$

$$= \left\{ \left( \left[ \frac{\partial^{2n} x}{\partial y^{2n}} \right]_{z=0} \right)^2 + \left( \left[ \frac{\partial^{2n} y}{\partial y^{2n}} \right]_{z=0} \right)^2 \right\}^{1/2}$$



$$\begin{aligned}
 &\leq \frac{4}{\pi} \frac{(2n)!}{a^{2n}} I_0\left(\frac{ac}{2}\right) \left\{ \max_{|z|=a} \left( e^{\frac{-cx}{2}} u \right)^2 + \max_{|z|=a} \left( e^{\frac{cx}{2}} v \right)^2 \right\}^{1/2} \\
 &\leq \frac{4}{\pi} \frac{(2n)!}{a^{2n}} I_0\left(\frac{ac}{2}\right) e^{\left|\frac{ca}{2}\right|} \left\{ [M(f)]^2 + [M(f)]^2 \right\}^{1/2} \\
 &= \frac{4}{\pi} 2^{1/2} \frac{(2n)!}{a^{2n}} M(f) e^{\left|\frac{ca}{2}\right|} I_0\left(\frac{ac}{2}\right) .
 \end{aligned}$$

Similarly we get the bound on  $|f^{[2n-1]}(0)|$  .

In the final result (3.5) we have used the fact that

$$\frac{2}{x} I_1(x) \leq I_0(x) . \quad \text{Q.E.D.}$$

COROLLARY 3.5: If  $f(z)$  is  $\Sigma$ -monogenic in  $|z-z_0| < a$  , then

$$|f^{[n]}(z_0)| \leq \frac{4}{\pi} 2^{1/2} \frac{M(f)}{a^n} n! e^{\left|\frac{ca}{2}\right| + |cx_0|} I_0\left(\frac{ac}{2}\right) .$$

From the bounds obtained on  $|a \cdot Z^{(n)}(0; z)|$  and  $|f^{[n]}(0)|$  the Generalized Taylor Theorem follows easily.

THEOREM 3.2: Suppose  $f(z)$  is  $\Sigma$ -monogenic in  $|z| < a$  , where

$$\Sigma = \begin{pmatrix} e^{-cx} & 1 \\ e^{-cx} & 1 \end{pmatrix} . \quad \text{Then}$$

$$f(z) = \sum_{n=0}^{\infty} \frac{f^{[n]}(0)}{n!} \cdot Z^{(n)}(0; z)$$

for all  $z$  such that  $|z| < a$  .



The above series converges absolutely for  $|z| < a$  and uniformly in any closed bounded circle  $|z| \leq a - \epsilon$ , ( $\epsilon > 0$ ).

PROOF: The statements on convergence follow immediately from Theorem 2.2 and equation (3.5).

$$\text{Thus } \sum_{n=0}^{\infty} \frac{f^{[n]}(0)}{n!} \cdot z^{(n)}(0; z) \text{ is } \Sigma\text{-monogenic for } |z| < a .$$

However, Bers and Gelbart have shown that

$$f(z) = \sum_{n=0}^{\infty} \frac{f^{[n]}(0)}{n!} z^{(n)}(0; z)$$

$$\text{for } |z| < \alpha < a . \quad \text{Therefore } \left\{ f(z) - \sum_{n=0}^{\infty} \frac{f^{[n]}(0)}{n!} \cdot z^{(n)}(0; z) \right\} \text{ is}$$

a function which is  $\Sigma$ -monogenic in  $|z| < a$  and equal to zero in some neighbourhood of the origin. Therefore it is identically zero in  $|z| < a$ .

Q.E.D.

COROLLARY 3.6: If  $f(z)$  is  $\Sigma$ -monogenic in  $|z - z_0| < a$ , then

$$f(z) = \sum_{n=0}^{\infty} \frac{f^{[n]}(z_0)}{n!} \cdot z^{(n)}(z_0; z) .$$

It is clear of course that the bound given for  $f^{[n]}(0)$  in Theorem 3.1 is not the best possible. The proof of Lemma 3.5 shows that the factor  $I_0(\frac{ca}{2})$  can be replaced by a factor which approaches one as  $n$  approaches infinity.





BIBLIOGRAPHY

1. L. Bers and A. Gelbart, On a Class of Functions Defined by Partial Differential Equations, Trans. Amer. Math. Soc. 56, 1944, 67-93.
2. L. Bers, The Expansion Theorem for Sigma-Monogenic Functions, Amer. J. Math., 72, 1950, 705-712.
3. H. Lamb, Hydrodynamics, Fifth Ed., Dover Publications, New York, 1924.
4. G.N. Watson, A Treatise on the Theory of Bessel Functions, 2nd Ed., Cambridge Univ. Press, London, 1958.
5. L. Bers and A. Gelbart, On a Class of Differential Equations in Mechanics of Continua, Quarterly of Applied Math., 1, No. 2, 1943, 168-188.



# APPENDIX

Theorem 1.2 can be extended to more general domains by a proof that merely uses integration by parts and not series expansions. We give the result here.

THEOREM A.1: Let  $f(z)$  be analytic in a domain each of whose points can be joined to the  $y$ -axis by a segment of a line parallel to the  $x$ -axis, such segment being interior to the domain. Then in such a domain  $\Gamma$ , define

$$i) \quad u_1 = f(y) + \frac{x}{\pi} \int_0^1 e^{cux} (1-u)^{1/2} u^{-1/2} du \int_0^\pi i \cos \phi f'(y+2ix[u(1-u)]^{1/2} \cos \phi) d\phi$$

$$v_1 = -\frac{x}{\pi} \int_0^1 e^{-cux} du \int_0^\pi f'(y+2ix[u(1-u)]^{1/2} \cos \phi) d\phi.$$

$$\begin{aligned} \text{Then,} \quad e^{-cx} u_x &= v_y \\ e^{-cx} u_y &= -v_x \quad \text{in } \Gamma. \end{aligned}$$

$$ii) \quad u_2 = \frac{x}{\pi} \int_0^1 e^{cux} du \int_0^\pi f'(y+2ix[u(1-u)]^{1/2} \cos \phi) d\phi$$

$$v_2 = f(y) + \frac{x}{\pi} \int_0^1 e^{-cux} (1-u)^{1/2} u^{-1/2} du \int_0^\pi i \cos \phi f'(y+2ix[u(1-u)]^{1/2} \cos \phi) d\phi.$$

$$\begin{aligned} \text{Then,} \quad e^{-cx} u_x &= v_y \\ e^{-cx} u_y &= -v_x \quad \text{in } \Gamma. \end{aligned}$$



PROOF: We need only prove i) since, regarding  $u_1$  and  $v_1$  as functions of  $c$ ,

$$\begin{aligned} u_2(c) &= -v_1(-c) \\ v_2(c) &= u_1(-c) . \end{aligned}$$

$$\begin{aligned} \text{Then,} \quad \frac{\partial u_1(c)}{\partial x} &= e^{cx} \frac{\partial v_1(c)}{\partial y} \\ \text{or} \quad \frac{\partial u_1(-c)}{\partial x} &= e^{-cx} \frac{\partial v_1(-c)}{\partial y} \\ \text{or} \quad \frac{\partial v_2(c)}{\partial x} &= -e^{-cx} \frac{\partial u_2(c)}{\partial y} \\ \text{or} \quad e^{-cx} \frac{\partial u_2(c)}{\partial x} &= - \frac{\partial v_2(c)}{\partial y} . \end{aligned}$$

$$\text{Similarly,} \quad e^{-cx} \frac{\partial u_2}{\partial x} = \frac{\partial v_2}{\partial y} .$$

We now proceed to prove i). (Call  $u_1 = u$ ,  $v_1 = v$ ).

$$\begin{aligned} v_y &= - \frac{x}{\pi} \int_0^1 e^{-cux} du \int_0^\pi f''(y+2ix[u(1-u)]^{1/2} \cos \varnothing) d\varnothing \\ u_x &= \frac{1}{\pi} \int_0^1 e^{cux} (1-u)^{1/2} u^{-1/2} du \int_0^\pi i \cos \varnothing f'(y+2ix[u(1-u)]^{1/2} \cos \varnothing) d\varnothing \\ &\quad - \frac{2x}{\pi} \int_0^1 e^{cux} (1-u) du \int_0^\pi \cos^2 \varnothing f''(y+2ix[u(1-u)]^{1/2} \cos \varnothing) d\varnothing \\ &\quad + \frac{cx}{\pi} \int_0^1 e^{cux} (1-u)^{1/2} u^{1/2} du \int_0^\pi i \cos \varnothing f'(y+2ix[u(1-u)]^{1/2} \cos \varnothing) d\varnothing . \end{aligned}$$



$$\begin{aligned}
 \text{So, } e^{-cx} u_x &= \frac{1}{\pi} \int_0^1 e^{-cxt} t^{1/2} (1-t)^{-1/2} dt \int_0^\pi i \cos \phi f'(y+2ix[t(1-t)])^{1/2} \cos \phi d\phi \\
 &\quad - \frac{2x}{\pi} \int_0^1 e^{-cxt} t dt \int_0^\pi \cos^2 \phi f''(y+2ix[t(1-t)])^{1/2} \cos \phi d\phi \\
 &\quad + \frac{cx}{\pi} \int_0^1 e^{-cxt} t^{1/2} (1-t)^{1/2} dt \int_0^\pi i \cos \phi f'(y+2ix[t(1-t)])^{1/2} \cos \phi d\phi \\
 &= F_1 + F_2 + F_3
 \end{aligned}$$

where we have made the substitution  $t = 1-u$ .

Now,

$$\begin{aligned}
 F_1 &= \frac{1}{\pi} \int_0^1 e^{-cxt} t^{1/2} (1-t)^{-1/2} dt \int_0^\pi f'(y+2ix[t(1-t)])^{1/2} \cos \phi d(i \sin \phi) \\
 &= - \frac{1}{\pi} \int_0^1 e^{-cxt} t^{1/2} (1-t)^{-1/2} dt \int_0^\pi i \sin \phi (-2ix[t(1-t)]^{1/2} \sin \phi) d\phi
 \end{aligned}$$

$$f''(y+2ix[t(1-t)])^{1/2} \cos \phi d\phi$$

[Integration by parts]

$$= - \frac{2x}{\pi} \int_0^1 e^{-cxt} t dt \int_0^\pi \sin^2 \phi f''(y+2ix[t(1-t)])^{1/2} \cos \phi d\phi$$

so that

$$F_1 + F_2 = - \frac{2x}{\pi} \int_0^1 e^{-cxt} t dt \int_0^\pi f''(y+2ix[t(1-t)])^{1/2} \cos \phi d\phi.$$





Next, integrating by parts in  $F_3$ , we get

$$\begin{aligned}
 F_3 &= -\frac{1}{\pi} \int_0^1 t^{1/2} (1-t)^{1/2} \left\{ \int_0^\pi i \cos \phi f'(y+2ix[t(1-t)]^{1/2} \cos \phi) d\phi \right\} d(e^{-cxt}) \\
 &= \frac{1}{\pi} \int_0^1 e^{-cxt} dt \left\{ \frac{1}{2} t^{-1/2} (1-t)^{-1/2} (1-2t) \int_0^\pi i \cos \phi f'(y+2ix[t(1-t)]^{1/2} \cos \phi) d\phi \right. \\
 &\quad \left. + t^{1/2} (1-t)^{1/2} \int_0^\pi (-x \cos^2 \phi) t^{-1/2} (1-t)^{-1/2} (1-2t) f''(y+2ix[t(1-t)]^{1/2} \cos \phi) d\phi \right\} \\
 &= \frac{1}{2\pi} \int_0^1 e^{-cxt} (1-2t) t^{-1/2} (1-t)^{-1/2} dt \int_0^\pi i \cos \phi f'(y+2ix[t(1-t)]^{1/2} \cos \phi) d\phi \\
 &\quad - \frac{x}{\pi} \int_0^1 (1-2t) e^{-cxt} dt \int_0^\pi \cos^2 \phi f''(y+2ix[t(1-t)]^{1/2} \cos \phi) d\phi \\
 &= -\frac{1}{2\pi} \int_0^1 e^{-cxt} (1-2t) t^{-1/2} (1-t)^{-1/2} dt \int_0^\pi (-i \sin \phi) (2ix[t(1-t)]^{1/2} \sin \phi) \cdot \\
 &\quad \cdot f''(y+2ix[t(1-t)]^{1/2} \cos \phi) d\phi \\
 &\quad - \frac{x}{\pi} \int_0^1 (1-2t) e^{-cxt} dt \int_0^\pi \cos^2 \phi f''(y+2ix[t(1-t)]^{1/2} \cos \phi) d\phi \\
 &= -\frac{x}{\pi} \int_0^1 e^{-cxt} (1-2t) dt \int_0^\pi f''(y+2ix[t(1-t)]^{1/2} \cos \phi) d\phi .
 \end{aligned}$$

$$\text{So } F_1 + F_2 + F_3 = -\frac{x}{\pi} \int_0^1 e^{-cxt} dt \int_0^\pi f''(y+2ix[t(1-t)]^{1/2} \cos \phi) d\phi .$$

$$\text{That is, } e^{-cx} u_x = v_y .$$

Now also,



$$u_y = f'(y) + \frac{x}{\pi} \int_0^1 e^{cux} (1-u)^{1/2} u^{-1/2} du \int_0^\pi i \cos \phi f''(y+2ix[u(1-u)])^{1/2} \cos \phi d\phi$$

$$\begin{aligned} v_x &= -\frac{1}{\pi} \int_0^1 e^{-cux} du \int_0^\pi f'(y+2ix[u(1-u)])^{1/2} \cos \phi d\phi \\ &\quad - \frac{x}{\pi} \int_0^1 e^{-cux} du \int_0^\pi 2i[u(1-u)]^{1/2} \cos \phi f''(y+2ix[u(1-u)])^{1/2} \cos \phi d\phi \\ &\quad + \frac{cx}{\pi} \int_0^1 e^{-cux} u du \int_0^\pi f'(y+2ix[u(1-u)])^{1/2} \cos \phi d\phi \end{aligned}$$

so that

$$\begin{aligned} e^{cx} v_x &= -\frac{2x}{\pi} \int_0^1 e^{cxt} t^{1/2} (1-t)^{1/2} dt \int_0^\pi i \cos \phi f''(y+2ix[t(1-t)])^{1/2} \cos \phi d\phi \\ &\quad - \frac{1}{\pi} \int_0^1 e^{cxt} dt \int_0^\pi f'(y+2ix[t(1-t)])^{1/2} \cos \phi d\phi \\ &\quad + \frac{cx}{\pi} \int_0^1 e^{cxt} (1-t) dt \int_0^\pi f'(y+2ix[t(1-t)])^{1/2} \cos \phi d\phi \\ &= G_1 + G_2 + G_3 . \end{aligned}$$

Now

$$\begin{aligned} G_3 &= \frac{1}{\pi} \int_0^1 (1-t) d(e^{cxt}) \int_0^\pi f'(y+2ix[t(1-t)])^{1/2} \cos \phi d\phi \\ &= \frac{1}{\pi} \left[ (1-t) \int_0^\pi f'(y+2ix[t(1-t)])^{1/2} \cos \phi d\phi e^{cxt} \right]_0^1 \\ &\quad + \frac{1}{\pi} \int_0^1 e^{cxt} dt \int_0^\pi f'(y+2ix[t(1-t)])^{1/2} \cos \phi d\phi \\ &\quad - \frac{1}{\pi} \int_0^1 e^{cxt} (1-t) dt \int_0^\pi 2ix \cos \phi \frac{1}{2} t^{-1/2} (1-t)^{-1/2} (1-2t) f''(y+2ix[t(1-t)])^{1/2} \cos \phi d\phi \end{aligned}$$



$$= -\frac{1}{\pi} \int_0^{\pi} f'(y) d\vartheta + \frac{1}{\pi} \int_0^1 e^{cxt} dt \int_0^{\pi} f'(y+2ix[t(1-t)]^{1/2} \cos \vartheta) d\vartheta \\ - \frac{x}{\pi} \int_0^1 e^{cxt} (1-t)^{1/2} t^{-1/2} dt \int_0^{\pi} (1-2t) i \cos \vartheta f''(y+2ix[t(1-t)]^{1/2} \cos \vartheta) d\vartheta$$

so that

$$G_3 + G_2 = -f'(y) - \frac{x}{\pi} \int_0^1 e^{cxt} (1-2t)(1-t)^{1/2} t^{-1/2} dt \int_0^{\pi} i \cos \vartheta \cdot \\ \cdot f''(y+2ix[t(1-t)]^{1/2} \cos \vartheta) d\vartheta .$$

Therefore,

$$G_1 + G_2 + G_3 = -f'(y) - \frac{x}{\pi} \int_0^1 e^{cxt} (1-t)^{1/2} t^{-1/2} dt \int_0^{\pi} i \cos \vartheta \cdot \\ \cdot f''(y+2ix[t(1-t)]^{1/2} \cos \vartheta) d\vartheta \\ = -u_y .$$

$$\text{Thus } e^{cx} v_x = -u_y \quad \text{or} \quad e^{-cx} u_y = -v_x .$$

The above procedure is valid wherever  $f(y+2ix[u(1-u)]^{1/2} \cos \vartheta)$  is analytic ( $0 \leq u \leq 1$ ,  $0 \leq \vartheta \leq \pi$ ). Since  $\max [u(1-u)]^{1/2} = \frac{1}{2}$ , the domain  $\Gamma$  will satisfy the condition. Q.E.D.

COROLLARY A.1: The formula given in Theorem 1.1 for the solutions of  $\Delta u - cu_x = 0$  and  $\Delta v + cv_x = 0$  are valid in  $\Gamma$ .













**B29871**